



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 02:08 PM UTC

PDB ID : 4BCA / pdb\_00004bca  
Title : MAMMALIAN ALKYLDIHYDROXYACETONEPHOSPHATE SYNTHASE: Tyr578Phe mutant  
Authors : Nenci, S.; Piano, V.; Rosati, S.; Aliverti, A.; Pandini, V.; Fraaije, M.W.; Heck, A.J.R.; Edmondson, D.E.; Mattevi, A.  
Deposited on : 2012-10-01  
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	658	<div> <div>%</div> <div> <div></div> <div>65%</div> <div>17%</div> <div>•</div> <div>16%</div> </div> </div>
1	B	658	<div> <div>2%</div> <div> <div></div> <div>64%</div> <div>17%</div> <div>•</div> <div>17%</div> </div> </div>
1	C	658	<div> <div>2%</div> <div> <div></div> <div>65%</div> <div>18%</div> <div>•</div> <div>15%</div> </div> </div>
1	D	658	<div> <div>%</div> <div> <div></div> <div>66%</div> <div>16%</div> <div>•</div> <div>16%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	CL	A	1659	-	-	X	-
4	SO4	B	1660	-	-	X	-

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 18202 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ALKYLDIHYDROXYACETONEPHOSPHATE SYNTHASE, PEROXISOMAL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	555	Total	C	N	O	S	0	0	0
			4393	2789	762	818	24			
1	B	543	Total	C	N	O	S	0	0	0
			4300	2732	747	797	24			
1	C	557	Total	C	N	O	S	0	2	0
			4412	2799	766	822	25			
1	D	550	Total	C	N	O	S	0	1	0
			4354	2761	757	811	25			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	578	PHE	TYR	engineered mutation	UNP P97275
B	578	PHE	TYR	engineered mutation	UNP P97275
C	578	PHE	TYR	engineered mutation	UNP P97275
D	578	PHE	TYR	engineered mutation	UNP P97275

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: C<sub>27</sub>H<sub>33</sub>N<sub>9</sub>O<sub>15</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	C	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	D	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		
3	B	1	Total	Cl	0	0
			1	1		
3	C	1	Total	Cl	0	0
			1	1		
3	D	1	Total	Cl	0	0
			1	1		

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	142	Total	O	0	0
			142	142		
5	B	95	Total	O	0	0
			95	95		
5	C	143	Total	O	0	0
			143	143		
5	D	112	Total	O	0	0
			112	112		







- Molecule 1: ALKYLDIHYDROXYACETONEPHOSPHATE SYNTHASE, PEROXISOMAL



- Molecule 1: ALKYLDIHYDROXYACETONEPHOSPHATE SYNTHASE, PEROXISOMAL







## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	61.68Å 99.32Å 107.98Å 90.43° 92.12° 95.20°	Depositor
Resolution (Å)	47.58 – 2.40 47.58 – 2.40	Depositor EDS
% Data completeness (in resolution range)	98.7 (47.58-2.40) 98.6 (47.58-2.40)	Depositor EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.32 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.187 , 0.252 0.190 , 0.250	Depositor DCC
$R_{free}$ test set	1091 reflections (1.09%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.8	Xtriage
Anisotropy	0.075	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 31.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.016 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	18202	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.30% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, FAD, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.13	2/4492 (0.0%)	1.16	9/6071 (0.1%)
1	B	1.07	1/4397 (0.0%)	1.14	13/5942 (0.2%)
1	C	1.16	8/4518 (0.2%)	1.16	10/6107 (0.2%)
1	D	1.12	5/4454 (0.1%)	1.14	13/6020 (0.2%)
All	All	1.12	16/17861 (0.1%)	1.15	45/24140 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	2

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	335	HIS	CG-ND1	-7.14	1.30	1.38
1	C	508	THR	CA-CB	7.07	1.65	1.53
1	C	335	HIS	CD2-NE2	-7.03	1.30	1.37
1	C	616	HIS	C-O	-6.72	1.17	1.24
1	C	507	LEU	C-O	-5.95	1.19	1.23

The worst 5 of 45 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	117	ARG	N-CA-C	8.42	120.54	111.36
1	B	298	THR	N-CA-C	-7.74	104.37	113.88
1	A	314	ILE	N-CA-C	-7.59	103.05	110.72
1	B	233	ILE	CA-C-N	-7.07	112.71	119.85

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	233	ILE	C-N-CA	-7.07	112.71	119.85

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	460	PRO	Peptide
1	C	507	LEU	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4393	0	4328	91	0
1	B	4300	0	4242	80	1
1	C	4412	0	4356	108	1
1	D	4354	0	4300	68	0
2	A	53	0	31	3	0
2	B	53	0	31	2	0
2	C	53	0	31	1	0
2	D	53	0	31	1	0
3	A	1	0	0	2	0
3	B	1	0	0	1	0
3	C	1	0	0	1	0
3	D	1	0	0	1	0
4	A	10	0	0	1	0
4	B	10	0	0	2	0
4	C	5	0	0	1	0
4	D	10	0	0	2	0
5	A	142	0	0	12	0
5	B	95	0	0	4	0
5	C	143	0	0	5	0
5	D	112	0	0	4	0
All	All	18202	0	17350	345	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 345 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:192:HIS:HB3	1:A:243:TYR:OH	1.37	1.25
1:C:508:THR:HA	1:C:511:ILE:CD1	1.65	1.24
1:A:421:MET:CG	5:A:2099:HOH:O	1.93	1.15
1:C:507:LEU:HD11	1:C:511:ILE:CG1	1.77	1.14
1:C:508:THR:HA	1:C:511:ILE:HD11	1.28	1.13

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:139:ASN:ND2	1:C:536:ARG:NH2[1_546]	2.14	0.06

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	549/658 (83%)	534 (97%)	15 (3%)	0	100	100
1	B	537/658 (82%)	514 (96%)	18 (3%)	5 (1%)	14	22
1	C	555/658 (84%)	528 (95%)	26 (5%)	1 (0%)	43	58
1	D	547/658 (83%)	525 (96%)	21 (4%)	1 (0%)	43	58
All	All	2188/2632 (83%)	2101 (96%)	80 (4%)	7 (0%)	36	50

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	504	GLY
1	B	195	PHE
1	B	196	LEU
1	C	506	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	503	ARG

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	476/545 (87%)	437 (92%)	39 (8%)	10	18
1	B	464/545 (85%)	432 (93%)	32 (7%)	14	24
1	C	479/545 (88%)	454 (95%)	25 (5%)	21	36
1	D	472/545 (87%)	438 (93%)	34 (7%)	13	23
All	All	1891/2180 (87%)	1761 (93%)	130 (7%)	14	24

5 of 130 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	424	GLN
1	D	482	LYS
1	B	195	PHE
1	B	180	ASP
1	D	507	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 52 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	656	ASN
1	C	424	GLN
1	D	597	GLN
1	C	110	GLN
1	C	388	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 4 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	SO4	B	1660	-	4,4,4	0.38	0	6,6,6	0.79	0
2	FAD	B	999	-	58,58,58	1.38	8 (13%)	85,89,89	1.94	25 (29%)
2	FAD	C	999	-	58,58,58	1.13	5 (8%)	85,89,89	1.99	25 (29%)
2	FAD	A	999	-	58,58,58	1.30	6 (10%)	85,89,89	1.66	16 (18%)
4	SO4	B	1661	-	4,4,4	0.31	0	6,6,6	0.72	0
4	SO4	D	1661	-	4,4,4	0.42	0	6,6,6	0.91	0
4	SO4	A	1660	-	4,4,4	0.23	0	6,6,6	1.01	0
4	SO4	C	1660	-	4,4,4	0.36	0	6,6,6	0.46	0
2	FAD	D	999	-	58,58,58	1.30	8 (13%)	85,89,89	1.76	24 (28%)
4	SO4	D	1660	-	4,4,4	0.17	0	6,6,6	0.49	0
4	SO4	A	1661	-	4,4,4	0.47	0	6,6,6	0.75	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	B	999	-	-	4/34/50/50	0/6/6/6
2	FAD	C	999	-	-	2/34/50/50	0/6/6/6

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	D	999	-	-	3/34/50/50	0/6/6/6
2	FAD	A	999	-	-	4/34/50/50	0/6/6/6

The worst 5 of 27 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	999	FAD	C4X-N5	6.13	1.44	1.30
2	D	999	FAD	C4X-N5	4.29	1.40	1.30
2	A	999	FAD	C4X-N5	4.03	1.39	1.30
2	D	999	FAD	PA-O3P	3.82	1.63	1.59
2	C	999	FAD	C4X-N5	3.67	1.38	1.30

The worst 5 of 90 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	999	FAD	C5A-C4A-N3A	-6.34	117.99	126.72
2	A	999	FAD	N3A-C2A-N1A	-5.91	119.63	128.58
2	C	999	FAD	N3A-C2A-N1A	-5.42	120.37	128.58
2	B	999	FAD	N9A-C8A-N7A	-5.21	106.55	113.94
2	B	999	FAD	C5A-N7A-C8A	4.57	110.64	103.45

There are no chirality outliers.

5 of 13 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	999	FAD	N10-C1'-C2'-O2'
2	A	999	FAD	N10-C1'-C2'-C3'
2	B	999	FAD	N10-C1'-C2'-O2'
2	B	999	FAD	N10-C1'-C2'-C3'
2	C	999	FAD	N10-C1'-C2'-O2'

There are no ring outliers.

9 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	1660	SO4	2	0
2	B	999	FAD	2	0
2	C	999	FAD	1	0
2	A	999	FAD	3	0
4	D	1661	SO4	1	0


*Continued on next page...*





*Continued from previous page...*


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	1660	SO4	1	0
2	D	999	FAD	1	0
4	D	1660	SO4	1	0
4	A	1661	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	555/658 (84%)	-0.40	8 (1%) 73 69	10, 24, 49, 73	0
1	B	543/658 (82%)	-0.21	12 (2%) 62 58	13, 30, 60, 75	0
1	C	557/658 (84%)	-0.41	11 (1%) 65 60	9, 24, 51, 73	2 (0%)
1	D	550/658 (83%)	-0.38	5 (0%) 81 78	9, 25, 53, 67	1 (0%)
All	All	2205/2632 (83%)	-0.35	36 (1%) 70 66	9, 25, 54, 75	3 (0%)

The worst 5 of 36 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	457	GLY	5.6
1	A	82	ILE	4.9
1	B	81	GLY	4.1
1	B	457	GLY	4.1
1	C	82	ILE	4.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	SO4	D	1661	5/5	0.87	0.13	41,44,53,53	0
4	SO4	B	1661	5/5	0.89	0.16	64,66,66,67	0
4	SO4	B	1660	5/5	0.91	0.14	47,49,51,51	0
3	CL	A	1659	1/1	0.92	0.12	36,36,36,36	0
3	CL	B	1659	1/1	0.93	0.16	56,56,56,56	0
4	SO4	A	1660	5/5	0.94	0.12	42,42,46,49	0
3	CL	C	1659	1/1	0.96	0.09	39,39,39,39	0
3	CL	D	1659	1/1	0.96	0.17	44,44,44,44	0
4	SO4	C	1660	5/5	0.97	0.09	44,44,47,49	0
2	FAD	A	999	53/53	0.98	0.05	9,11,14,16	0
4	SO4	A	1661	5/5	0.98	0.06	30,30,33,33	0
4	SO4	D	1660	5/5	0.98	0.09	36,38,39,43	0
2	FAD	B	999	53/53	0.98	0.05	12,15,18,21	0
2	FAD	D	999	53/53	0.99	0.04	8,10,13,15	0
2	FAD	C	999	53/53	0.99	0.04	7,9,12,15	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around FAD B 999:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around FAD D 999:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers [i](#)

There are no such residues in this entry.