



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 18, 2026 – 05:34 AM UTC

PDB ID : 6CPU / pdb\_00006cpu  
Title : Crystal structure of yeast caPDE2  
Authors : Ke, H.; Wang, y.  
Deposited on : 2018-03-14  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	571	<div> <div>16%</div> <div>89%</div> <div>8%</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4504 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphodiesterase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	526	Total	C	N	O	S	0	0	0
			4271	2763	702	788	18			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	316	LYS	THR	conflict	UNP Q8NJP9

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	231	Total	O	0	0
			231	231		



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	139.52Å 74.75Å 65.46Å 90.00° 109.12° 90.00°	Depositor
Resolution (Å)	65.91 – 1.80 65.91 – 1.80	Depositor EDS
% Data completeness (in resolution range)	98.9 (65.91-1.80) 98.9 (65.91-1.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.75 (at 1.81Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, $R_{free}$	0.199 , 0.226 0.208 , 0.235	Depositor DCC
$R_{free}$ test set	2826 reflections (4.83%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.9	Xtriage
Anisotropy	0.020	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 33.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4504	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.19% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.52	0/4362	0.74	0/5917

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4271	0	4297	9	0
2	A	1	0	0	0	0
3	A	1	0	0	0	0
4	A	231	0	0	0	0
All	All	4504	0	4297	9	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (9) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:134:TRP:CZ3	1:A:268:LYS:HD3	2.39	0.58
1:A:524:ASP:HA	1:A:526:HIS:CE1	2.41	0.56
1:A:134:TRP:CD2	1:A:268:LYS:HE3	2.46	0.51
1:A:290:LEU:HD13	1:A:339:THR:HG23	1.92	0.50
1:A:523:PRO:O	1:A:524:ASP:CG	2.58	0.47
1:A:268:LYS:HD2	1:A:353:HIS:O	2.15	0.47
1:A:524:ASP:C	1:A:524:ASP:OD2	2.60	0.43
1:A:139:HIS:HA	1:A:142:TYR:CZ	2.54	0.43
1:A:175:ARG:HD2	1:A:183:LEU:HD12	2.01	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	516/571 (90%)	503 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	485/528 (92%)	478 (99%)	7 (1%)	59	52

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	104	GLN
1	A	262	ILE
1	A	342	LEU
1	A	367	PRO
1	A	429	GLU
1	A	524	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	99	ASN
1	A	104	GLN
1	A	139	HIS
1	A	170	ASN
1	A	182	GLN
1	A	224	ASN
1	A	244	GLN
1	A	317	HIS
1	A	388	ASN
1	A	428	ASN
1	A	465	ASN
1	A	526	HIS
1	A	561	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.



There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	526/571 (92%)	0.86	92 (17%) 4 3	16, 30, 62, 102	1 (0%)

All (92) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	443	ILE	6.7
1	A	569	ALA	6.3
1	A	546	ASP	5.9
1	A	142	TYR	5.3
1	A	52	THR	4.7
1	A	530	ILE	4.5
1	A	439	THR	4.3
1	A	140	PHE	4.3
1	A	435	LYS	4.3
1	A	531	PHE	4.2
1	A	438	LYS	4.1
1	A	521	ILE	4.0
1	A	559	MET	4.0
1	A	76	THR	3.9
1	A	109	ASP	3.9
1	A	535	LEU	3.9
1	A	563	LEU	3.9
1	A	570	LYS	3.9
1	A	528	GLY	3.8
1	A	245	VAL	3.8
1	A	247	GLY	3.7
1	A	434	LEU	3.7
1	A	324	GLN	3.6
1	A	111	GLY	3.6
1	A	323	LEU	3.6
1	A	514	GLU	3.5
1	A	12	ILE	3.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	567	GLU	3.3
1	A	566	LEU	3.2
1	A	560	GLU	3.2
1	A	442	GLU	3.2
1	A	539	ASN	3.2
1	A	525	ILE	3.2
1	A	445	ASN	3.2
1	A	440	HIS	3.1
1	A	436	SER	3.1
1	A	520	GLU	3.1
1	A	139	HIS	2.9
1	A	524	ASP	2.9
1	A	9	ASP	2.8
1	A	556	ASP	2.8
1	A	13	PRO	2.8
1	A	545	SER	2.8
1	A	519	LEU	2.7
1	A	547	LEU	2.7
1	A	555	CYS	2.7
1	A	516	ARG	2.7
1	A	431	VAL	2.7
1	A	330	GLU	2.7
1	A	406	LYS	2.6
1	A	549	PRO	2.6
1	A	526	HIS	2.6
1	A	568	ARG	2.6
1	A	544	VAL	2.6
1	A	430	TYR	2.5
1	A	437	PHE	2.5
1	A	515	LEU	2.5
1	A	557	ILE	2.5
1	A	428	ASN	2.5
1	A	104	GLN	2.5
1	A	548	LEU	2.5
1	A	432	ASN	2.5
1	A	564	PHE	2.5
1	A	484	PHE	2.4
1	A	14	GLN	2.4
1	A	404	GLU	2.4
1	A	537	ALA	2.4
1	A	444	LEU	2.4
1	A	118	SER	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	171	CYS	2.4
1	A	534	ASN	2.3
1	A	322	ALA	2.3
1	A	10	LEU	2.3
1	A	248	PHE	2.2
1	A	337	ILE	2.2
1	A	518	ILE	2.2
1	A	540	LEU	2.2
1	A	532	PHE	2.1
1	A	542	ASN	2.1
1	A	565	TRP	2.1
1	A	113	LEU	2.1
1	A	110	GLU	2.1
1	A	1	MET	2.1
1	A	75	ASN	2.1
1	A	441	ASN	2.1
1	A	359	ASP	2.1
1	A	527	LYS	2.0
1	A	541	PHE	2.0
1	A	156	SER	2.0
1	A	114	TYR	2.0
1	A	523	PRO	2.0
1	A	303	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
-----	------	-------	-----	-------	------	-----	-----------------------------	-------

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	MG	A	602	1/1	0.97	0.06	26,26,26,26	0
2	ZN	A	601	1/1	0.98	0.12	34,34,34,34	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.