



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 02:47 PM UTC

PDB ID : 7DEV / pdb\_00007dev  
Title : Crystal Structures of Anthocyanin 5,3'-aromatic acyltransferase from *Gentiana triflora*  
Authors : Murayama, K.; Kato-Murayama, M.; Shirouzu, M.  
Deposited on : 2020-11-05  
Resolution : 3.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1456 (3.10-3.10)
Clashscore	190562	1539 (3.10-3.10)
Ramachandran outliers	187476	1467 (3.10-3.10)
Sidechain outliers	187428	1467 (3.10-3.10)
RSRZ outliers	180081	1456 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	476	

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 3601 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Anthocyanin 5-aromatic acyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	454	3601	2334	586	662	19	0	0	0


There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	GLY	-	expression tag	UNP Q9ZWR8
A	-5	SER	-	expression tag	UNP Q9ZWR8
A	-4	PHE	-	expression tag	UNP Q9ZWR8
A	-3	THR	-	expression tag	UNP Q9ZWR8
A	-2	SER	-	expression tag	UNP Q9ZWR8
A	-1	SER	-	expression tag	UNP Q9ZWR8
A	0	GLY	-	expression tag	UNP Q9ZWR8

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Anthocyanin 5-aromatic acyltransferase



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	51.44Å 112.99Å 183.97Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.54 – 3.10 49.54 – 3.10	Depositor EDS
% Data completeness (in resolution range)	99.8 (49.54-3.10) 99.8 (49.54-3.10)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.23	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.42 (at 3.12Å)	Xtriage
Refinement program	PHENIX 1.13_2998, CNS	Depositor
R, $R_{free}$	0.191 , 0.251 0.193 , 0.254	Depositor DCC
$R_{free}$ test set	1017 reflections (9.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	40.8	Xtriage
Anisotropy	1.224	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 66.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	3601	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.54% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.38	0/3691	0.60	0/5006

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3601	0	3602	97	0
All	All	3601	0	3602	97	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (97) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:73:THR:HG22	1:A:192:ILE:HD12	1.66	0.76
1:A:73:THR:HG21	1:A:189:TRP:HA	1.66	0.76
1:A:368:GLU:HA	1:A:372:HIS:HB2	1.68	0.76
1:A:289:VAL:HG11	1:A:431:ILE:HD13	1.68	0.76
1:A:192:ILE:HD11	1:A:201:LEU:HD11	1.70	0.73

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:364:GLY:O	1:A:367:ILE:HG13	1.94	0.66
1:A:436:SER:HA	1:A:444:GLU:HG3	1.78	0.66
1:A:5:GLN:HG2	1:A:7:VAL:H	1.63	0.63
1:A:142:ILE:HG23	1:A:152:PRO:HG3	1.81	0.63
1:A:263:LEU:HB3	1:A:284:MET:HE2	1.82	0.62
1:A:432:TYR:CZ	1:A:446:GLY:HA3	2.35	0.62
1:A:295:LYS:HE2	1:A:351:VAL:HA	1.83	0.61
1:A:266:LYS:HG2	1:A:356:LEU:HD23	1.82	0.61
1:A:202:LEU:HB2	1:A:207:LEU:HD22	1.82	0.60
1:A:138:MET:HE3	1:A:155:ALA:HB2	1.84	0.58
1:A:125:LEU:HG	1:A:249:VAL:HG21	1.86	0.57
1:A:378:LEU:HA	1:A:381:ALA:HB2	1.85	0.57
1:A:69:SER:O	1:A:73:THR:HG23	2.05	0.57
1:A:77:TYR:OH	1:A:184:MET:HG2	2.04	0.56
1:A:365:GLU:HG2	1:A:369:LYS:HE3	1.88	0.56
1:A:116:ASP:CG	1:A:164:ARG:HH12	2.13	0.55
1:A:269:ASN:C	1:A:271:ARG:H	2.14	0.55
1:A:257:LEU:HD22	1:A:441:LYS:HB2	1.87	0.55
1:A:399:ILE:HA	1:A:431:ILE:O	2.06	0.55
1:A:321:CYS:SG	1:A:339:ALA:HB2	2.48	0.54
1:A:28:LEU:HB3	1:A:96:PHE:HB2	1.90	0.53
1:A:277:ILE:HD13	1:A:361:ALA:HB2	1.89	0.53
1:A:182:PHE:O	1:A:186:ILE:HG13	2.09	0.53
1:A:36:PRO:HA	1:A:229:MET:HE3	1.91	0.52
1:A:310:ASN:N	1:A:310:ASN:OD1	2.43	0.52
1:A:138:MET:HE2	1:A:152:PRO:HB2	1.92	0.51
1:A:201:LEU:HD12	1:A:201:LEU:H	1.75	0.51
1:A:417:PRO:O	1:A:437:ARG:NH1	2.45	0.50
1:A:319:ALA:HB1	1:A:371:LEU:HD12	1.94	0.49
1:A:67:LYS:HG2	1:A:108:LEU:HD13	1.93	0.49
1:A:12:LYS:HE3	1:A:107:THR:HG21	1.95	0.49
1:A:356:LEU:O	1:A:360:VAL:HG23	2.12	0.49
1:A:193:ASN:HD22	1:A:412:PHE:HB3	1.77	0.49
1:A:250:ARG:HH11	1:A:250:ARG:HG3	1.78	0.48
1:A:188:ALA:HB1	1:A:201:LEU:HD23	1.95	0.48
1:A:428:ALA:HB3	1:A:430:LEU:HG	1.95	0.48
1:A:231:ASP:O	1:A:235:MET:HG3	2.13	0.48
1:A:290:TRP:O	1:A:294:VAL:HG23	2.13	0.48
1:A:77:TYR:HB3	1:A:80:LEU:HD12	1.97	0.47
1:A:178:ASP:HB2	1:A:338:LEU:HD11	1.96	0.47
1:A:252:THR:HG21	1:A:419:LYS:HE2	1.97	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:153:LEU:HD13	1:A:176:ILE:HG12	1.96	0.47
1:A:432:TYR:CE2	1:A:446:GLY:HA3	2.49	0.47
1:A:63:ILE:HD12	1:A:110:VAL:HG22	1.97	0.47
1:A:211:ASP:OD1	1:A:213:SER:OG	2.31	0.47
1:A:173:HIS:O	1:A:176:ILE:HG22	2.15	0.46
1:A:192:ILE:HG12	1:A:199:ALA:HB3	1.97	0.46
1:A:233:LEU:O	1:A:237:SER:HB3	2.14	0.46
1:A:117:PHE:HB3	1:A:164:ARG:HD3	1.98	0.46
1:A:51:PHE:CZ	1:A:53:TYR:HD2	2.34	0.46
1:A:346:THR:HG22	1:A:348:LYS:N	2.31	0.46
1:A:113:SER:HB3	1:A:161:PHE:CE2	2.51	0.45
1:A:407:SER:HB3	1:A:420:PHE:CD2	2.51	0.45
1:A:381:ALA:C	1:A:383:THR:H	2.24	0.45
1:A:116:ASP:HB3	1:A:119:TYR:HB3	1.99	0.44
1:A:324:LEU:HD11	1:A:371:LEU:CD2	2.47	0.44
1:A:30:VAL:HG21	1:A:94:PRO:HB2	2.00	0.44
1:A:30:VAL:HG23	1:A:95:LYS:HA	1.99	0.44
1:A:190:ALA:HA	1:A:412:PHE:CD2	2.52	0.44
1:A:18:PRO:HD3	1:A:71:SER:O	2.17	0.44
1:A:117:PHE:H	1:A:164:ARG:HH11	1.65	0.44
1:A:289:VAL:HG21	1:A:445:ILE:HG21	2.01	0.43
1:A:373:ASN:C	1:A:375:LYS:H	2.26	0.43
1:A:383:THR:HG22	1:A:387:GLU:HG3	2.00	0.43
1:A:170:LEU:HD12	1:A:170:LEU:N	2.34	0.43
1:A:138:MET:CE	1:A:152:PRO:HB2	2.49	0.43
1:A:402:SER:OG	1:A:404:LYS:HG2	2.17	0.43
1:A:293:MET:HG2	1:A:397:LEU:HD13	1.99	0.43
1:A:326:THR:HA	1:A:327:PRO:C	2.44	0.43
1:A:72:LEU:O	1:A:75:LYS:HG2	2.18	0.42
1:A:449:LEU:HB2	1:A:454:MET:HE2	2.01	0.42
1:A:18:PRO:HB2	1:A:21:THR:OG1	2.19	0.42
1:A:242:LYS:HE3	1:A:242:LYS:HB3	1.81	0.42
1:A:354:LYS:HD3	1:A:357:LEU:HD23	2.01	0.42
1:A:402:SER:HB3	1:A:405:PHE:CE2	2.55	0.42
1:A:330:PRO:HD2	1:A:333:TYR:HB2	2.01	0.42
1:A:420:PHE:C	1:A:420:PHE:CD1	2.98	0.41
1:A:51:PHE:CZ	1:A:62:VAL:HG21	2.55	0.41
1:A:370:ARG:HH22	1:A:387:GLU:CD	2.28	0.41
1:A:317:PHE:CD1	1:A:317:PHE:N	2.89	0.41
1:A:51:PHE:HB3	1:A:165:GLY:HA2	2.03	0.41
1:A:153:LEU:O	1:A:172:ALA:HA	2.21	0.41

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:309:GLU:HB3	1:A:310:ASN:H	1.52	0.41
1:A:347:HIS:O	1:A:351:VAL:HG22	2.20	0.41
1:A:28:LEU:HD12	1:A:28:LEU:HA	1.84	0.41
1:A:86:MET:HA	1:A:87:PRO:HD3	1.94	0.41
1:A:142:ILE:CG2	1:A:152:PRO:HG3	2.49	0.41
1:A:312:LEU:HD23	1:A:312:LEU:HA	1.91	0.41
1:A:271:ARG:HD3	1:A:271:ARG:HA	1.85	0.41
1:A:145:MET:HE3	1:A:145:MET:HB2	1.83	0.40
1:A:153:LEU:HD23	1:A:153:LEU:HA	1.78	0.40
1:A:260:ILE:HD13	1:A:281:THR:HG23	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	450/476 (94%)	421 (94%)	26 (6%)	3 (1%)	18 49

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	270	LEU
1	A	274	GLU
1	A	145	MET

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	400/420 (95%)	381 (95%)	19 (5%)	23	54

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	13	CYS
1	A	148	TYR
1	A	149	LYS
1	A	214	ILE
1	A	216	LYS
1	A	231	ASP
1	A	234	GLU
1	A	242	LYS
1	A	249	VAL
1	A	276	THR
1	A	309	GLU
1	A	310	ASN
1	A	317	PHE
1	A	326	THR
1	A	377	VAL
1	A	386	SER
1	A	414	TRP
1	A	424	SER
1	A	448	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	14	GLN
1	A	97	GLN
1	A	193	ASN
1	A	230	GLN
1	A	265	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	454/476 (95%)	-0.08	10 (2%) 62 41	13, 34, 83, 129	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	374	GLU	3.2
1	A	301	VAL	3.1
1	A	20	ASP	2.7
1	A	318	THR	2.7
1	A	389	ASN	2.7
1	A	452	ILE	2.4
1	A	60	ASP	2.3
1	A	371	LEU	2.2
1	A	164	ARG	2.1
1	A	278	ARG	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.