



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 02:48 AM UTC

PDB ID : 4EIW / pdb\_00004eiw  
Title : Whole cytosolic region of atp-dependent metalloprotease FtsH (G399L)  
Authors : Suno, R.; Niwa, H.; Tsuchiya, D.; Yoshida, M.; Morikawa, K.  
Deposited on : 2012-04-06  
Resolution : 3.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1270 (4.10-3.70)
Clashscore	190562	1034 (4.08-3.72)
Ramachandran outliers	187476	1251 (4.10-3.70)
Sidechain outliers	187428	1243 (4.10-3.70)
RSRZ outliers	180081	1269 (4.10-3.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	508	<div> <div>14%</div> <div>48%</div> <div>26%</div> <div>•</div> <div>10%</div> </div>
1	B	508	<div> <div>8%</div> <div>48%</div> <div>26%</div> <div>5%</div> <div>12%</div> </div>
1	C	508	<div> <div>14%</div> <div>48%</div> <div>25%</div> <div>•</div> <div>10%</div> </div>
1	D	508	<div> <div>8%</div> <div>50%</div> <div>25%</div> <div>5%</div> <div>12%</div> </div>
1	E	508	<div> <div>15%</div> <div>47%</div> <div>26%</div> <div>•</div> <div>10%</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	508	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ADP	B	2001	-	-	X	-
2	ADP	C	1001	-	-	X	-
2	ADP	D	2001	-	-	X	-
2	ADP	E	1001	-	-	X	-
2	ADP	F	2001	-	-	X	-

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 21429 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP-dependent zinc metalloprotease FtsH.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	458	Total	C	N	O	S	0	0	0
			3578	2245	658	662	13			
1	B	446	Total	C	N	O	S	0	0	0
			3511	2206	641	651	13			
1	C	458	Total	C	N	O	S	0	0	0
			3578	2245	658	662	13			
1	D	446	Total	C	N	O	S	0	0	0
			3511	2206	641	651	13			
1	E	458	Total	C	N	O	S	0	0	0
			3578	2245	658	662	13			
1	F	446	Total	C	N	O	S	0	0	0
			3511	2206	641	651	13			

There are 60 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	117	GLY	-	expression tag	UNP Q5SI82
A	118	PRO	-	expression tag	UNP Q5SI82
A	119	LEU	-	expression tag	UNP Q5SI82
A	120	GLY	-	expression tag	UNP Q5SI82
A	121	SER	-	expression tag	UNP Q5SI82
A	122	HIS	-	expression tag	UNP Q5SI82
A	123	MET	-	expression tag	UNP Q5SI82
A	124	GLY	-	expression tag	UNP Q5SI82
A	125	ALA	-	expression tag	UNP Q5SI82
A	399	LEU	GLY	engineered mutation	UNP Q5SI82
B	117	GLY	-	expression tag	UNP Q5SI82
B	118	PRO	-	expression tag	UNP Q5SI82
B	119	LEU	-	expression tag	UNP Q5SI82
B	120	GLY	-	expression tag	UNP Q5SI82
B	121	SER	-	expression tag	UNP Q5SI82
B	122	HIS	-	expression tag	UNP Q5SI82
B	123	MET	-	expression tag	UNP Q5SI82

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
B	124	GLY	-	expression tag	UNP Q5SI82
B	125	ALA	-	expression tag	UNP Q5SI82
B	399	LEU	GLY	engineered mutation	UNP Q5SI82
C	117	GLY	-	expression tag	UNP Q5SI82
C	118	PRO	-	expression tag	UNP Q5SI82
C	119	LEU	-	expression tag	UNP Q5SI82
C	120	GLY	-	expression tag	UNP Q5SI82
C	121	SER	-	expression tag	UNP Q5SI82
C	122	HIS	-	expression tag	UNP Q5SI82
C	123	MET	-	expression tag	UNP Q5SI82
C	124	GLY	-	expression tag	UNP Q5SI82
C	125	ALA	-	expression tag	UNP Q5SI82
C	399	LEU	GLY	engineered mutation	UNP Q5SI82
D	117	GLY	-	expression tag	UNP Q5SI82
D	118	PRO	-	expression tag	UNP Q5SI82
D	119	LEU	-	expression tag	UNP Q5SI82
D	120	GLY	-	expression tag	UNP Q5SI82
D	121	SER	-	expression tag	UNP Q5SI82
D	122	HIS	-	expression tag	UNP Q5SI82
D	123	MET	-	expression tag	UNP Q5SI82
D	124	GLY	-	expression tag	UNP Q5SI82
D	125	ALA	-	expression tag	UNP Q5SI82
D	399	LEU	GLY	engineered mutation	UNP Q5SI82
E	117	GLY	-	expression tag	UNP Q5SI82
E	118	PRO	-	expression tag	UNP Q5SI82
E	119	LEU	-	expression tag	UNP Q5SI82
E	120	GLY	-	expression tag	UNP Q5SI82
E	121	SER	-	expression tag	UNP Q5SI82
E	122	HIS	-	expression tag	UNP Q5SI82
E	123	MET	-	expression tag	UNP Q5SI82
E	124	GLY	-	expression tag	UNP Q5SI82
E	125	ALA	-	expression tag	UNP Q5SI82
E	399	LEU	GLY	engineered mutation	UNP Q5SI82
F	117	GLY	-	expression tag	UNP Q5SI82
F	118	PRO	-	expression tag	UNP Q5SI82
F	119	LEU	-	expression tag	UNP Q5SI82
F	120	GLY	-	expression tag	UNP Q5SI82
F	121	SER	-	expression tag	UNP Q5SI82
F	122	HIS	-	expression tag	UNP Q5SI82
F	123	MET	-	expression tag	UNP Q5SI82
F	124	GLY	-	expression tag	UNP Q5SI82
F	125	ALA	-	expression tag	UNP Q5SI82

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	399	LEU	GLY	engineered mutation	UNP Q5SI82


- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: C<sub>10</sub>H<sub>15</sub>N<sub>5</sub>O<sub>10</sub>P<sub>2</sub>).






- Molecule 1: ATP-dependent zinc metalloprotease FtsH






• Molecule 1: ATP-dependent zinc metalloprotease FtsH

Chain D: 8% 50% 25% 5% 12%



• Molecule 1: ATP-dependent zinc metalloprotease FtsH

Chain E: 15% 47% 26% 10%



[illegible]

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	146.15Å 146.15Å 349.06Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	71.53 – 3.90 71.53 – 3.90	Depositor EDS
% Data completeness (in resolution range)	97.1 (71.53-3.90) 97.1 (71.53-3.90)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	7.46 (at 3.89Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.299 , 0.312 0.298 , 0.309	Depositor DCC
$R_{free}$ test set	1967 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	99.1	Xtriage
Anisotropy	0.272	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 568.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.27$ , $\langle L^2 \rangle = 0.11$	Xtriage
Estimated twinning fraction	0.237 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.81	EDS
Total number of atoms	21429	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	112.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.77% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.80	3/3636 (0.1%)	1.17	22/4906 (0.4%)
1	B	0.82	2/3568 (0.1%)	1.26	40/4815 (0.8%)
1	C	0.82	3/3636 (0.1%)	1.22	34/4906 (0.7%)
1	D	0.81	2/3568 (0.1%)	1.24	35/4815 (0.7%)
1	E	0.81	8/3636 (0.2%)	1.19	25/4906 (0.5%)
1	F	0.78	0/3568	1.23	32/4815 (0.7%)
All	All	0.80	18/21612 (0.1%)	1.22	188/29163 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	5
1	C	0	1
1	D	0	6
1	E	0	1
1	F	0	5
All	All	0	20

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	214	ARG	CZ-NH2	-8.43	1.22	1.33
1	C	214	ARG	CZ-NH2	7.85	1.43	1.33
1	B	586	GLU	CD-OE1	-6.81	1.12	1.25
1	A	214	ARG	CZ-NH2	-6.71	1.24	1.33
1	E	319	ARG	CZ-NH1	-6.65	1.23	1.32

The worst 5 of 188 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	511	HIS	CA-C-N	-12.31	104.45	119.84
1	D	511	HIS	C-N-CA	-12.31	104.45	119.84
1	F	511	HIS	CA-C-N	-11.79	105.11	119.84
1	F	511	HIS	C-N-CA	-11.79	105.11	119.84
1	A	177	ASN	CB-CA-C	-11.37	98.74	110.33

There are no chirality outliers.

5 of 20 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	213	ALA	Peptide
1	A	532	VAL	Peptide
1	B	244	ALA	Peptide
1	B	288	GLY	Peptide
1	B	382	ARG	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3578	0	3623	901	5
1	B	3511	0	3556	894	3
1	C	3578	0	3623	843	0
1	D	3511	0	3556	892	3
1	E	3578	0	3623	844	3
1	F	3511	0	3556	896	2
2	A	27	0	12	8	0
2	B	27	0	12	9	0
2	C	27	0	12	10	0
2	D	27	0	12	9	0
2	E	27	0	12	11	0
2	F	27	0	12	10	0
All	All	21429	0	21609	5119	8

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 119.

The worst 5 of 5119 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:416:ALA:CB	1:E:577:LEU:HD23	1.18	1.63
1:A:416:ALA:HB3	1:A:577:LEU:CD2	1.33	1.58
1:A:416:ALA:CB	1:A:577:LEU:HD23	1.15	1.55
1:F:376:ALA:CA	1:F:381:ARG:HD2	1.31	1.55
1:E:416:ALA:HB3	1:E:577:LEU:CD2	1.35	1.55

The worst 5 of 8 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:570:GLU:CG	1:D:382:ARG:NH2[3_564]	1.61	0.59
1:A:417:TYR:OH	1:D:382:ARG:NE[3_564]	1.74	0.46
1:A:177:ASN:OD1	1:E:214:ARG:NH2[6_665]	1.85	0.35
1:B:238:ARG:NE	1:F:378:ARG:NH2[6_665]	1.91	0.29
1:A:570:GLU:CB	1:D:382:ARG:NH2[3_564]	2.08	0.12

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	456/508 (90%)	323 (71%)	111 (24%)	22 (5%)	2	18
1	B	442/508 (87%)	289 (65%)	125 (28%)	28 (6%)	1	15
1	C	456/508 (90%)	323 (71%)	112 (25%)	21 (5%)	2	19
1	D	442/508 (87%)	288 (65%)	121 (27%)	33 (8%)	1	12
1	E	456/508 (90%)	324 (71%)	110 (24%)	22 (5%)	2	18
1	F	442/508 (87%)	288 (65%)	120 (27%)	34 (8%)	1	12
All	All	2694/3048 (88%)	1835 (68%)	699 (26%)	160 (6%)	1	15

5 of 160 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	153	VAL
1	A	511	HIS
1	B	153	VAL
1	B	274	GLU
1	B	379	GLU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	365/402 (91%)	229 (63%)	136 (37%)	0	1
1	B	361/402 (90%)	224 (62%)	137 (38%)	0	1
1	C	365/402 (91%)	225 (62%)	140 (38%)	0	1
1	D	361/402 (90%)	222 (62%)	139 (38%)	0	1
1	E	365/402 (91%)	228 (62%)	137 (38%)	0	1
1	F	361/402 (90%)	225 (62%)	136 (38%)	0	1
All	All	2178/2412 (90%)	1353 (62%)	825 (38%)	0	1

5 of 825 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	285	GLU
1	E	215	VAL
1	F	503	ARG
1	D	321	ILE
1	D	284	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 36 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	204	HIS
1	F	511	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	F	277	GLN
1	F	333	GLN
1	B	431	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	ADP	D	2001	-	28,29,29	1.54	6 (21%)	43,45,45	2.03	10 (23%)
2	ADP	B	2001	-	28,29,29	1.53	5 (17%)	43,45,45	1.87	11 (25%)
2	ADP	F	2001	-	28,29,29	1.54	5 (17%)	43,45,45	1.94	12 (27%)
2	ADP	E	1001	-	28,29,29	1.47	4 (14%)	43,45,45	1.87	9 (20%)
2	ADP	A	1001	-	28,29,29	1.52	4 (14%)	43,45,45	1.93	10 (23%)
2	ADP	C	1001	-	28,29,29	1.51	4 (14%)	43,45,45	2.07	10 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the



Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.  
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	D	2001	-	-	2/16/32/32	0/3/3/3
2	ADP	B	2001	-	-	3/16/32/32	0/3/3/3
2	ADP	F	2001	-	-	2/16/32/32	0/3/3/3
2	ADP	E	1001	-	-	2/16/32/32	0/3/3/3
2	ADP	A	1001	-	-	1/16/32/32	0/3/3/3
2	ADP	C	1001	-	-	2/16/32/32	0/3/3/3

The worst 5 of 28 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1001	ADP	C5-C4	5.16	1.48	1.39
2	F	2001	ADP	C5-C4	5.12	1.48	1.39
2	E	1001	ADP	C5-C4	5.08	1.48	1.39
2	C	1001	ADP	C5-C4	4.92	1.47	1.39
2	B	2001	ADP	C5-C4	4.88	1.47	1.39

The worst 5 of 62 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	2001	ADP	C5-C4-N3	-6.87	117.26	126.72
2	C	1001	ADP	C5-C4-N3	-6.36	117.95	126.72
2	A	1001	ADP	C5-C4-N3	-6.11	118.30	126.72
2	E	1001	ADP	C5-C4-N3	-5.92	118.56	126.72
2	F	2001	ADP	C5-C4-N3	-5.86	118.64	126.72

There are no chirality outliers.

5 of 12 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1001	ADP	C5'-O5'-PA-O1A
2	B	2001	ADP	C5'-O5'-PA-O2A
2	C	1001	ADP	C5'-O5'-PA-O1A
2	C	1001	ADP	C5'-O5'-PA-O3A
2	D	2001	ADP	C5'-O5'-PA-O3A



There are no ring outliers.



6 monomers are involved in 57 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	2001	ADP	9	0
2	B	2001	ADP	9	0
2	F	2001	ADP	10	0
2	E	1001	ADP	11	0
2	A	1001	ADP	8	0
2	C	1001	ADP	10	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	458/508 (90%)	-0.31	1 (0%) 91 81	21, 118, 174, 229	0
1	B	446/508 (87%)	-0.29	1 (0%) 91 81	16, 125, 176, 241	0
1	C	458/508 (90%)	-0.21	0 100 100	21, 129, 184, 237	0
1	D	446/508 (87%)	-0.28	2 (0%) 88 74	23, 103, 173, 241	0
1	E	458/508 (90%)	-0.22	1 (0%) 91 81	24, 130, 179, 223	0
1	F	446/508 (87%)	-0.23	1 (0%) 91 81	21, 128, 173, 261	0
All	All	2712/3048 (88%)	-0.25	6 (0%) 91 81	16, 124, 177, 261	0

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	600	GLU	2.8
1	A	528	GLY	2.4
1	B	382	ARG	2.4
1	D	443	ARG	2.3
1	E	270	GLY	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	ADP	B	2001	27/27	0.97	0.06	65,72,75,79	0
2	ADP	C	1001	27/27	0.97	0.07	70,73,74,76	0
2	ADP	F	2001	27/27	0.97	0.07	64,69,74,74	0
2	ADP	E	1001	27/27	0.98	0.07	66,70,72,73	0
2	ADP	A	1001	27/27	0.98	0.06	63,66,70,71	0
2	ADP	D	2001	27/27	0.99	0.07	58,64,74,74	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around ADP C 1001:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around ADP F 2001:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around ADP E 1001:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around ADP A 1001:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around ADP D 2001:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.